



Unexpectedly long hospital stays as an indicator of risk of unsafe care. Dutch experiences with a new outcome indicator.

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2013-004773
Article Type:	Research
Date Submitted by the Author:	30-Dec-2013
Complete List of Authors:	Borghans, Ine; Dutch Healthcare Inspectorate (IGZ), Hekkert, Karin; Dutch Hospital Data, den Ouden, Lya; Dutch Healthcare Inspectorate (IGZ), Cihangir, Sezgin; Dutch Hospital Data, ; Vesseur, Jan; Dutch Healthcare Inspectorate (IGZ), Kool, Rudolf; Radboud university medical centre, Scientific Institute for Quality of Healthcare (IQ Healthcare Westert, Gert; Radboud university medical centre, Scientific Institute for Quality of Healthcare (IQ Healthcare
Primary Subject Heading:	Health services research
Secondary Subject Heading:	Health policy
Keywords:	Health & safety < HEALTH SERVICES ADMINISTRATION & MANAGEMENT, Risk management < HEALTH SERVICES ADMINISTRATION & MANAGEMENT, Quality in health care < HEALTH SERVICES ADMINISTRATION & MANAGEMENT

SCHOLARONE™
Manuscripts

Unexpectedly long hospital stays as an indicator of risk of unsafe care.

Dutch experiences with a new outcome indicator.

Ine Borghans^{1,2§}, Karin D Hekkert³, Lya den Ouden², Sezgin Cihangir³, Jan Vesseur², Rudolf B Kool¹, Gert P Westert¹

¹ Scientific Institute for Quality of Healthcare (IQ Healthcare), Radboud university medical centre, P.O. Box 9101, 6500 HB, Nijmegen, the Netherlands

- Ine Borghans, Senior research fellow
- Rudolf B Kool, Senior research fellow
- Gert P Westert, Head of the department

² Dutch Health Care Inspectorate (IGZ), P.O. Box 2680, 3500 GR, Utrecht, the Netherlands

- Ine Borghans, Coordinator Risk Analyzes
- Lya den Ouden, Coordinating specialistic Inspector
- Jan Vesseur, Project Chief Inspector Patient Safety

³ Dutch Hospital Data (DHD), P.O. Box 9696, 3506 GR Utrecht, the Netherlands

- Karin D Hekkert, Researcher
- Sezgin Cihangir, Researcher

[§] Corresponding author

Tijn Kool, P.O. Box 9101, 6500 HB, Nijmegen, the Netherlands

T: +31(0)611531143

E: tijn.kool@radboudumc.nl

Keywords: quality indicator, adverse events, length of stay, patient safety, hospital care

Word count : 2941

ABSTRACT

Background: We developed an outcome indicator based on the finding that complications often prolong the patient's hospital stay. A higher percentage of patients with an unexpectedly long length of stay (UL-LOS) compared to the national average may indicate shortcomings in the patient safety. Since 2009, the Dutch Health Care Inspectorate has added the UL-LOS indicator to their supervision framework. In this article we report the Dutch experiences with the UL-LOS and some of its characteristics.

Methods: The indicator is calculated among survivors and based on the percentage of patients with a prolonged length of stay of more than 50% of the expected length of stay. In order to compare hospitals properly we used data of standardised length of stay for the years 2008 - 2011. The standardisation was based on patients' age, primary diagnosis and main procedure. We explored the utility of the UL-LOS indicator by measuring its variability across hospitals and the stability over time. We also examined the correlation between the UL-LOS and standardised mortality. Finally, we investigated whether the indicator was influenced by hospitals that did have problems discharging their patients to other health services such as nursing homes.

Results: The UL-LOS indicator showed considerable variability between the Dutch hospitals: from 8.6% to 20.1% in 2011. The outcomes had relatively small confidence intervals since they were based on large numbers of patients. The stability of the indicator over time was quite high. The indicator had a significant positive correlation with the standardised mortality, and no significant correlation with the percentage of patients that was discharged to other facilities than other hospitals and home.

Conclusions: The UL-LOS indicator is a useful addition to other patient safety indicators by revealing variation between hospitals and areas of possible patient safety improvement.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

STRENGTHS AND LIMITATIONS OF THIS STUDY


- This study provides a hospital-wide indicator that can be used in addition to mortality and readmissions rates in order to identify potential safety risks
- The current indicator adjusts for differences in age, principal diagnosis and procedures. But there are probably more variables involved in a prolonged hospital stay.
- The indicator currently ‘counts’ all patients from whom the actual LOS exceeds the expected duration by 50% or more. It needs to be studied whether this cut-off point can be set separately for each patient group.

For peer review only

BACKGROUND

For about ten years, improving quality of care based on outcome indicators is seen as an essential component in optimising safety in healthcare. In the Netherlands, like in many other countries, a large number of indicators have been developed and introduced to monitor the quality and safety of hospital care[1-3]. Many of these indicators concern a specific patient group. There are also some general quality indicators that concern the whole hospital. The most important ones are: unexpectedly long length of stay (UL-LOS); unplanned readmissions and higher than expected mortality (measured by the Hospital Standardised Mortality Rate, HSMR) as shown in Figure 1.

Figure 1. General outcome indicators which may reveal poor or sub-optimal quality of hospital care



According to the Dutch Healthcare Inspectorate (DHI) these are especially of interest for identifying general patient safety risks in hospital care. It is important to measure all three indicators since a degree of substitution –or competition- between them is possible. For example, if a hospital tries to discharge patients too quickly, this policy could result in a higher percentage of unplanned readmissions. The indicator for unplanned readmissions, already used in several countries such as the UK and USA[4-7], is not yet available in the Netherlands. The HSMR has already been available in Dutch hospitals since 2006[8].

The UL-LOS did not exist until some years ago the DHI was looking for more general patient safety indicators. Research had shown that hospital adverse events often result in a longer length of stay[9-12][13][14, 15][16-19][20, 21]. In several other studies on adverse events, a long length of stay was used as an important trigger for selecting medical records[22-24]. So, if complications often prolong the patient's hospital stay, could an outcome indicator be developed such as the percentage of patients with an unexpectedly long length of stay (UL-LOS) compared to the national average? Such an outcome may indicate shortcomings in the quality or safety of care delivered by the hospital.

As far back as 1999 Silber et al. had already published research about an indicator called ‘conditional length of stay’. This was based on LOS-data and took into account the fact that patient stays tend to become prolonged after complications. They developed this indicator by testing if LOS distributions display an ‘extended’ pattern of decreasing hazards after a transition point. This would suggest that ‘the longer a patient has stayed in the hospital, the longer a patient will likely stay in the hospital’. Or, alternatively, there is the possibility that ‘the longer a patient has stayed in the hospital, the faster a patient will likely be discharged from the hospital’[25].

Based on these former ideas, the DHI decided to introduce such an indicator of prolonged length of stay. In the current paper we explore the utility of this indicator by measuring the variability of the indicator across hospitals and the stability over time. We also examine the correlation between the two existing indicators of the model: the UL-LOS and the HSMR, as they are both supposed to be an indicator of risk of unsafe care. The HSMR focuses by definition on adverse events leading to hospital mortality. The UL-LOS is calculated among survivors and involves a much wider range of adverse events. It includes all adverse events leading to a substantial prolonged length of stay. We expected a positive relationship between the two indicators, as reduced quality of care leads to more adverse events, and more adverse events lead to more patients with prolonged hospitalisation as well as to more deaths.

METHODS


Definition

First, we defined an outcome indicator that suited the purpose of identifying adverse events. We used former research (Ref Silber and Bottle and Aylin) to choose a single cut-off percentage prolonged length of stay across all diagnoses and case mixes to distinguish between ‘normal’ variation in LOS and variation in LOS that might be caused by complications and other patient safety issues[26]. We formulated the indicator ‘Unexpectedly long length of stay’ as the percentage of clinically admitted patients with an actual hospital stay that was more than 50% longer than expected[2]. We excluded from this indicator patients who died in hospital.

We used a threshold of 50% for three reasons:

1. We especially wanted to include patients that stayed longer because of complications and adverse events, and not patients that just stayed a little bit longer in hospital because of variations in efficiency, see figure 2. The percentage should not be too small.

Figure 2: components of LOS



2. In literature we found the use of a threshold of 75% [27, 28], which in fact was arbitrary chosen.
3. We analysed for the Netherlands the percentages of patients with a longer than expected LOS for ten different threshold groups between 0 and 100%.

Table 1. All Dutch hospital patients divided by LOS-groups.

		Percentage of patients 2009	Percentage of patients 2010	Percentage of patients 2011
UL-LOS	observed LOS >100% longer than expected LOS	8,1%	7,9%	7,1%
	observed LOS 90-100% longer than expected LOS	0,9%	0,9%	0,8%
	observed LOS 80-90% longer than expected LOS	1,2%	1,0%	0,9%
	observed LOS 70-80% longer than expected LOS	1,1%	1,1%	1,0%
	observed LOS 60-70% longer than expected LOS	1,2%	1,3%	1,2%
	observed LOS 50-60% longer than expected LOS	1,6%	1,7%	1,6%
NON UL-LOS	observed LOS 40-50% longer than expected LOS	2,2%	2,0%	1,9%
	observed LOS 30-40% longer than expected LOS	2,1%	2,2%	2,0%
	observed LOS 20-30% longer than expected LOS	3,1%	2,9%	2,7%
	observed LOS 10-20% longer than expected LOS	3,2%	3,5%	3,4%
	observed LOS <10% longer than expected LOS	4,6%	5,0%	4,8%
	observed LOS ≤ expected LOS	68,6%	68,5%	70,8%
	deceased patients	2,1%	2,0%	1,8%
TOTAL		100,0%	100,0%	100,0%

We found that the percentages after the threshold of 50% seem to dip beneath 2%.

In combination with our first experiences with case studies that showed that the number of adverse events clearly increased from a longer than expected length of stay of 50% or more[26], we decided to use the threshold of 50% in order not to exclude too many patients with adverse events. In fact the threshold depends to what specificity or what false positive rates you are willing to accept. With a threshold of 50% we expect to find a higher number of adverse events than with a threshold of 75%, but we expect the proportion of adverse events will be lower. A higher threshold would mean more efficiency in finding adverse events.

Data

Second, we looked for data that were routinely registered for administrative purposes. Using an existing registration minimalizes extra registration burden.

These databases can be used to predict risks with similar discrimination to clinical databases[29]. The National Medical Registration (LMR) already exists for 50 years in the Netherlands and contains data of hospital admissions including medical data such as diagnosis and surgical procedures as well as patient-specific data such as age and hospital stay[30]. The LMR also includes the variable ‘expected length of stay’, which is generated by indirect standardisation based on the following three patient characteristics, which are the most important characteristics for standardisation of length of stay data[31]:

- Age: Divided into 5 categories: 0, 1-14, 15-44, 45-64, 65+ years;
- Primary diagnosis: This is the main diagnosis that led to the admission; it includes about 1,000 diagnoses classified by the ICD9 in three digits;
- Morbidity group: Morbidity groups mainly divide patients with or without procedures. For patients with procedures the morbidity groups are made by unique combinations of a diagnosis and one or more procedures. On average it includes five procedure groups. Procedures are classified by the Dutch Classification System of Procedures.

Together these three parameters produced $5 \times 1,000 \times 5 = 25,000$ cells for which the mean length of stay is taken as the expected length of stay.

Sample

Third, we selected hospitals for which the indicator could be calculated. Hospitals had to participate fully in the LMR, not only by the registration of the diagnoses of clinical patients but also of the procedures. For our study this resulted in the use of data of the year 2011 of 61 out of all 90 Dutch hospitals. 6 hospitals were excluded because they did not participate at all, 5 hospitals were excluded because they participated for less than 50% and 18 hospitals were excluded because they did not register the procedures in the LMR. Among these hospitals were general and tertiary teaching hospitals, and they did not differ in size or region from the hospitals that could be included in the study.

To optimize the comparability of hospitals, we stratified the sample into three groups: 32 general hospitals, 21 tertiary teaching hospitals (TTHs) and 8 university medical centres (UMCs). Within these groups, size, degree of specialisation, financing system and complexity of patients are comparable[32-34].

Analyses

Fourth, to explore the utility of the indicator, we measured the variability of the indicator across hospitals and the stability over time. To find out whether the indicator is stable over time, we determined the correlation between the percentages per hospital in '2008 and 2009', '2009 and 2010', '2010 and 2011' and '2008 and 2011'.

Fifth, in order to analyse whether the indicator could identify risks of unsafe care, we correlated the results of the UL-LOS with the HSMRs for the year 2011. Therefore we calculated the Pearson correlations and the 2-tailed Sig. between the unexpectedly long lengths of stay and HSMRs. The HSMR consists of the quotients of observed mortality and expected mortality in 50 diagnostic groups (CCS) in which 80% of all hospital mortality took place. The expected mortality was based on the following characteristics of the patients: age, sex, CCS-subgroup, co morbidity (Charlson index), urgency, social deprivation, source organisation type, month and year.

As with the UL-LOS, the HSMR could only be calculated for hospitals that participated in the LMR. Some additional criteria were used in order to optimize reliability of the standardisation. To be included in the HSMR, hospitals had to:

- Avoid the use of vague diagnostic codes (this had to be less than 2% of the admissions);
- Perform an adequate registration of the urgency of the admission (more than 30% of the admissions had to be marked as urgent);
- Perform an adequate registration of the comorbidity of patients (the mean number of secondary diagnosis per admission had to be more than 0,5).

In addition, the HSMR had to count for more than 70% of all deaths in hospital, and the hospital had to have more than 50 expected deaths per year. All these additional criteria resulted in 58 hospitals remaining for our correlation study between UL-LOS and HSMR for the year 2011.

Sixth, we investigated whether the UL-LOS indicator (and the HSMR) might be influenced by hospitals that, for local resource reasons, do not have as much access to nursing home beds, home care support, palliative care or hospices, and thus keep more of these patients in the hospital rather than discharging them to stay or die elsewhere. We addressed this by two different analyses:


1) we calculated the percentage of patients that was discharged to other facilities than other hospitals and home. This percentage approximately gives the extent to which hospitals are able to discharge patients for long-term or palliative care.

2) We re-ran the analysis restricted to diagnostic categories where most patients are discharged home without a diagnosis for which one might expect a palliative approach. To select this group, we excluded all CCS groups[35] with carcinoma.

RESULTS

In Figure 3 the percentage of unexpectedly long length of stay is given for the 61 hospitals in our study. The figure also shows the 95% confidence limits. A distinction has been made between general hospitals, TTHs and UMCs. The national median of the percentage of clinical admissions with a UL-LOS was 11.3%. The UMCs had a relatively high score on this indicator compared with the TTHs and general hospitals.

Figure 3. Percentage of admissions with an unexpectedly long length of stay (UL-LOS) for the hospitals in our study, defined by type of hospital: general hospitals (N=32), tertiary teaching hospitals (TTHs) (N=21) and University Medical Centres (UMCs) (N=8); LMR 2011



For the UMCs the variation of the percentages was between 12.0% and 20.1%, with a median of 15.1%. The TTHs varied between 9.4% and 15.0%, with a median of 11.3%. For the general hospitals the variation was between 8.6% and 16.0%, with a median of 11.1%. With an independent samples t-test we found no significant difference between tertiary teaching hospitals and general hospitals, $t(53) = 0.16$; $p = .88$.

To explore the stability of the indicator, we calculated the correlation of the indicator between two subsequent years. The R^2 between 2008 and 2009 was 0.89, between 2009 and 2010 it was 0.86 and between 2010 and 2011 it was 0.90. Figure 4 shows the correlation between 2008 and 2011. The R^2 between these years was 0.70.

Figure 4. Correlation between the years 2008 and 2011 for the percentage of admissions with an unexpectedly long length of stay (UL-LOS) per hospital. LMR 2008 and 2011 (N=57)




Table 2 shows for the year 2011 for each hospital the UL-LOS indicator and the HSMR in two ways: with and without patients with carcinoma. This table also shows for each hospital the percentage of patients admitted to other destinations than home or other hospital. There was no substantial difference in the outcome of the UL-LOS indicator calculated with or without patients with carcinomas. The correlation of the UL-LOS with/without carcinomas was 1,00; correlation of the HSMR with/without carcinomas was 0,96. The Pearson correlation between the UL-LOS indicator and the HSMR with all diagnosis groups was 0,44 ($p < 0.001$) and with carcinoma excluded 0,52 ($p < 0.001$). We found no significant correlation between the UL-LOS indicator and the score on discharging patients. The Pearson correlation between the UL-LOS and discharging palliative patients with all diagnosis groups was -0.15 ($p < 0.001$) and with carcinoma excluded -0.16 ($p < 0.001$).

Table 2. 61 hospitals with their UL-LOS and HSMR in two ways: with and without patients with carcinoma and the percentage of patients admitted to other destinations than home or other hospital; 2011.

	Total hospital		Carcinoma excluded		
Hospital number	UL-LOS 2011	HSMR 2011	UL-LOS 2011	HSMR 2011	admissions to other destinations than home or other hospital
1	20,1%	117	20,3%	128	1,4%
2	18,7%	not available	18,8%	not available	0,6%
3	17,8%	115	17,8%	118	0,5%
4	16,0%	147	15,5%	145	3,4%
5	15,6%	118	15,6%	128	1,8%
6	15,6%	129	15,8%	133	1,4%
7	15,0%	100	15,0%	101	4,7%
8	14,7%	101	14,5%	103	0,9%
9	14,7%	not available	14,6%	not available	2,5%
10	14,7%	101	14,6%	100	4,3%
11	13,9%	93	13,6%	98	1,7%
12	13,6%	108	13,4%	105	3,2%
13	13,4%	113	13,3%	109	2,3%
14	13,4%	106	13,6%	104	not available
15	13,2%	71	13,0%	70	2,2%
16	13,0%	107	12,9%	112	1,2%
17	12,8%	112	12,6%	112	4,5%
18	12,7%	100	12,7%	101	1,7%
19	12,4%	108	12,2%	107	2,2%
20	12,4%	65	12,1%	64	3,1%
21	12,4%	106	12,2%	102	2,3%
22	12,3%	92	12,1%	88	2,1%
23	12,2%	70	12,1%	72	1,4%
24	12,1%	94	12,1%	86	2,1%
25	12,0%	91	12,3%	96	2,7%
26	11,7%	104	11,6%	102	0,2%
27	11,5%	78	11,5%	82	3,4%
28	11,5%	116	11,4%	111	1,2%
29	11,4%	115	11,5%	114	2,6%
30	11,4%	101	11,1%	105	1,0%
31	11,3%	101	11,0%	90	0,8%
32	11,3%	94	11,3%	96	2,8%
33	11,3%	94	11,3%	98	2,8%
34	11,3%	107	11,3%	115	2,3%
35	11,2%	91	11,2%	91	3,3%

36	11,2%	83	11,1%	80	0,6%
37	11,1%	104	11,3%	106	1,5%
38	11,1%	114	10,8%	111	1,8%
39	11,0%	103	10,9%	108	2,4%
40	11,0%	93	10,7%	91	2,2%
41	10,9%	114	10,9%	114	2,4%
42	10,9%	106	10,8%	103	2,9%
43	10,9%	107	10,8%	105	2,0%
44	10,9%	104	10,5%	101	2,5%
45	10,7%	110	10,6%	101	3,4%
46	10,7%	81	10,6%	79	2,4%
47	10,6%	100	10,4%	99	1,6%
48	10,6%	99	10,6%	95	2,4%
49	10,6%	not available	10,2%	not available	2,2%
50	10,5%	93	10,3%	95	2,3%
51	10,4%	91	10,1%	88	3,1%
52	10,1%	107	10,0%	103	3,0%
53	10,1%	81	10,4%	78	2,0%
54	9,9%	101	9,7%	104	3,1%
55	9,9%	61	9,6%	65	3,8%
56	9,7%	111	9,7%	112	1,2%
57	9,6%	99	9,5%	101	2,9%
58	9,4%	82	9,5%	85	3,1%
59	9,4%	83	9,2%	77	1,3%
60	9,3%	89	9,3%	86	2,7%
61	8,6%	91	8,7%	93	1,0%

DISCUSSION

In this paper, we described the development of a patient safety indicator for Dutch hospitals that, to the best of our knowledge, has not been described in the literature till now. The indicator is defined as the percentage of clinically admitted patients with an actual hospital stay that was more than 50% longer than expected. The indicator showed considerable variability between the Dutch hospitals: from 8.6% to 20.1% in 2011. It also showed serious variation within homogenous groups of hospitals. The stability of the indicator over three years was quite high and the indicator had a significant positive correlation with the HSMR. This indicates that the hospitals with more patients with a UL-LOS were also the hospitals with higher standardised mortality. This might support our hypothesis that sub-optimal quality of care may both lead to more patients with an unexpectedly long length of stay as well as to more mortality than expected. More research – especially record reviewing to look for adverse events - is needed for a stronger support of this hypothesis.

The strong correlation between the UL-LOS with and without carcinomas, and the low correlation between UL-LOS and discharging patients, seem to indicate that the UL-LOS is not influenced by differences between hospitals in the ability to admit patients to next care facilities. This might be important for hospitals in realizing that high percentages of patients with an ULLOS will probably not be caused by opportunities of discharging patients.

In terms of evaluation of care it is becoming increasingly common for hospitals to study patient records retrospectively, especially in cases of deceased patients^{25,26}. This indicator might be a good research tool to identify records of patients who were discharged alive. This could lead to opportunities for improvements that are different from those based on patient record reviewing after death.

The indicator provides insight into the percentage of patients that stayed at least 50% longer than expected. The assumption is that in this group a relatively large number of patients have had to deal with unexpected developments in their disease, resulting in complications that cause a prolonged stay. It could be much more effective for hospitals to review records of hospital admissions selected by this indicator compared to randomly selected patient records. Reviewing records takes considerable time and by using this indicator for selection, time could be saved by reviewing fewer records from which more lessons might be learnt[26].

Hospital management might also have financial reasons to be interested in the indicator in addition to the quality and safety aspects of the UL-LOS indicator. Having a longer than expected length of stay, costs money in the present Dutch financing system. Although the indicator is not developed for financial purposes, the use of the indicator may, as a beneficial side effect, also reduce the total amount of hospital days.

There are two main limitations to this study.

1. Some hospitals, especially the UMCs, differed from the national median. The high score for the UMCs could indicate that there is insufficient adjustment for the specific patient categories admitted to the UMCs. The current indicator adjusts for differences in age, principal diagnosis and procedures. But there are probably more variables involved in a prolonged hospital stay. The case mix adjustment is more limited than for example the HSMR. Further research is needed to determine which other patient characteristics, e.g. co-morbidity, play a significant role in a prolonged length of stay and whether they are of added value in standardising LOS next to the present co-founders.
2. We used a threshold of a length of stay of 50% longer than expected for the indicator. This threshold was based on the three abovementioned reasons and especially our first experiences with reviewing hospital records based on LOS[26]. However, there is no

evidence that the threshold has to be exactly 50%. A more detailed study is needed to determine the appropriateness of this threshold. Further research is also needed to determine to what extent the proportion of patients which crosses the threshold can vary for each combination of age, main diagnosis and procedure. If the variations between hospitals are large, and the case mix clearly differs, this will present a number of unequal opportunities that will result in crossing the 50% threshold. We might need to vary the threshold for different diagnostic groups.

CONCLUSION

The DHI introduced the outcome indicator 'Percentage of patients with an unexpectedly long LOS' for its supervision since 2010. It is based on the assumption that complications often prolong the patient's hospital stay. A higher percentage of patients with a UL-LOS compared to the national average, after a correction is made for case mix variations, may indicate shortcomings in the quality or safety of care delivered by the hospital. The indicator currently 'counts' all patients from whom the actual LOS exceeds the expected duration by 50% or more. It needs to be studied whether this cut-off point can be set separately for each patient group. The indicator varies systematically between hospitals and is rather stable over time. It correlates with other outcome indicators, which could indicate the capacity to identify opportunities for improvement of patient safety.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ACKNOWLEDGMENTS

FUNDING

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

CONTRIBUTORSHIP STATEMENT

IB designed the study, performed the analysis, interpreted the results and drafted the manuscript. KDH and SC contributed to the analysis and the interpretation of the results. RBK, LdO and JV helped to interpret the results and contributed to the discussion. RBK and GPW supervised the study and participated in the formulation of the discussion. All authors reviewed and edited the manuscript for intellectual content. All authors read and approved the final manuscript.

COMPETING INTERESTS

The authors declare that they have no competing interests.

DATA SHARING STATEMENT

Not relevant for this study

REFERENCES

- 1 Inspectie voor de Gezondheidszorg [Dutch Health Care Inspectorate]. *Basisset kwaliteitsindicatoren ziekenhuizen 2013*. Utrecht: IGZ; 2012.
- 2 Inspectie voor de Gezondheidszorg [Dutch Health Care Inspectorate]. *Basisset veiligheidsindicatoren ziekenhuizen 2010-2012*. Utrecht: IGZ; 2010.
- 3 Kwaliteitsindicatoren Zichtbare Zorg Ziekenhuizen. Available at: <http://www.zichtbarezorg.nl/page/Ziekenhuizen-en-ZBC-s/Kwaliteitsindicatoren#ZI>. Accessed 12/01, 2011.
- 4 Nolte E, Roland M, Guthrie S, et al. Preventing emergency readmissions to hospital, A scoping review. RAND Europe; 2012.
- 5 Lagoe RJ, Nanno DS, Luziani ME. Quantitative tools for addressing hospital readmissions. *BMC Res Notes* 2012;**5**:620,0500-5-620.
- 6 Goldfield NI, McCullough EC, Hughes JS, et al. Identifying potentially preventable readmissions. *Health Care Financ Rev* 2008;**30**:75-91.
- 7 Joynt KE, Jha AK. Thirty-day readmissions--truth and consequences. *N Engl J Med* 2012;**366**:1366-9.
- 8 Jarman B, Pieter D, van der Veen A, et al. The hospital standardised mortality ratio: a powerful tool for Dutch hospitals to assess their quality of care?. *Qual Saf Health Care* 2010;**19**:9-13.
- 9 Hoonhout LH, de Bruijne MC, Wagner C, et al. Nature, occurrence and consequences of medication-related adverse events during hospitalization: a retrospective chart review in the Netherlands. *Drug Saf* 2010;**33**:853-64.
- 10 Hoonhout LH, de Bruijne MC, Wagner C, et al. Direct medical costs of adverse events in Dutch hospitals. *BMC Health Serv Res* 2009;**9**:27.
- 11 Sari AB, Sheldon TA, Cracknell A, et al. Extent, nature and consequences of adverse events: results of a retrospective casenote review in a large NHS hospital. *Qual Saf Health Care* 2007;**16**:434-9.

12 Ehsani JP, Jackson T, Duckett SJ. The incidence and cost of adverse events in Victorian hospitals 2003-04. *Med J Aust* 2006;**184**:551-5.

13 Cho SH, Ketefian S, Barkauskas VH, et al. The effects of nurse staffing on adverse events, morbidity, mortality, and medical costs. *Nurs Res* 2003;**52**:71-9.

14 Camp M, Chang DC, Zhang Y, et al. Risk factors and outcomes for foreign body left during a procedure: analysis of 413 incidents after 1 946 831 operations in children. *Arch Surg* 2010;**145**:1085-90.

15 Lotfipour S, Kaku SK, Vaca FE, et al. Factors associated with complications in older adults with isolated blunt chest trauma. *West J Emerg Med* 2009;**10**:79-84.

16 Williams DJ, Olsen S, Crichton W, et al. Detection of adverse events in a Scottish hospital using a consensus-based methodology. *Scott Med J* 2008;**53**:26-30.

17 Kaushal R, Bates DW, Franz C, et al. Costs of adverse events in intensive care units. *Crit Care Med* 2007;**35**:2479-83.

18 Rice-Townsend S, Hall M, Jenkins KJ, et al. Analysis of adverse events in pediatric surgery using criteria validated from the adult population: justifying the need for pediatric-focused outcome measures. *J Pediatr Surg* 2010;**45**:1126-36.

19 Schioler T, Lipczak H, Pedersen BL, et al. Incidence of adverse events in hospitals. A retrospective study of medical records. *Ugeskr Laeger* 2001;**163**:5370-8.

20 Baker GR, Norton PG, Flintoft V, et al. The Canadian Adverse Events Study: the incidence of adverse events among hospital patients in Canada. *CMAJ* 2004;**170**:1678-86.

21 Davis P, Lay-Yee R, Briant R, et al. Adverse events in New Zealand public hospitals I: occurrence and impact. *N Z Med J* 2002;**115**:U271.

- 22 de Bruijne MC, Zegers M, Hoonhout LHF, et al. Onbedoelde schade in Nederlandse ziekenhuizen: dossieronderzoek van ziekenhuisopnames in 2004. Amsterdam: EMGO Instituut/VUmc en Nivel, Nederlands Instituut voor onderzoek van de gezondheidszorg; 2007.
- 23 Langelaan M, Baines RJ, Broekens MA, et al. Monitor Zorggerelateerde Schade 2008, dossieronderzoek in Nederlandse ziekenhuizen. NIVEL; 2010.
- 24 Zegers M, de Bruijne MC, Wagner C, et al. Adverse events and potentially preventable deaths in Dutch hospitals: results of a retrospective patient record review study. *Qual Saf Health Care* 2009;**18**:297-302.
- 25 Silber JH, Rosenbaum PR, Koziol LF, et al. Conditional Length of Stay. *Health Serv Res* 1999;**34**:349-63.
- 26 Cihangir S, Borghans I, Hekkert K, et al. A pilot study on record reviewing with a priori patient selection. *BMJ Open* 2013;**3**:10.1136/bmjopen.2013-003034. Print 2013.
- 27 Bottle A, Aylin P. Intelligent information: a national system for monitoring clinical performance. *Health Serv Res* 2008;**43**:10-31.
- 28 Bottle A, Middleton S, Kalkman CJ, et al. Global comparators project: international comparison of hospital outcomes using administrative data. *Health Serv Res* 2013;**48**:2081-100.
- 29 Aylin P, Bottle A, Majeed A. Use of administrative data or clinical databases as predictors of risk of death in hospital: comparison of models. *BMJ* 2007;**334**:1044.
- 30 Landelijke Medische Registratie (LMR). Available at: <http://www.dutchhospitaldata.nl/registraties/lmrlazr/Paginas/default.aspx>. Accessed 06/10, 2013.
- 31 Hart 't P, de Boer K, Draaisma D. On the length of stay in hospitals. Radboud University Nijmegen; 2012.
- 32 Website of more than 40 small and medium-sized Dutch hospitals. They take more than a third of all Dutch hospital care on their behalf. Available at: <http://www.saz-ziekenhuizen.nl>. Accessed 09-30, 2013.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

33 Website with facts and figures of all (8) Dutch university hospitals. Available at: <http://www.nfu.nl/umc/feiten/>. Accessed 09-30, 2013.

34 Website of 28 Dutch tertiary teaching hospitals. Available at: <http://www.stz.nl/pagina/9-over-stz.html>. Accessed 09-30, 2013.

35 Rivard PE, Luther SL, Christiansen CL, et al. Using patient safety indicators to estimate the impact of potential adverse events on outcomes. *Med Care Res Rev* 2008;**65**:67-87.

BMJ Open

Unexpectedly long hospital stays as an indicator of risk of unsafe care: an exploratory study.

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2013-004773.R1
Article Type:	Research
Date Submitted by the Author:	15-Apr-2014
Complete List of Authors:	Borghans, Ine; Dutch Healthcare Inspectorate (IGZ), Hekkert, Karin; Dutch Hospital Data, den Ouden, Lya; Dutch Healthcare Inspectorate (IGZ), Cihangir, Sezgin; Dutch Hospital Data, Vesseur, Jan; Dutch Healthcare Inspectorate (IGZ), Kool, Rudolf; Radboud university medical centre, Scientific Institute for Quality of Healthcare (IQ healthcare) Westert, Gert; Radboud university medical centre, Scientific Institute for Quality of Healthcare (IQ healthcare)
Primary Subject Heading:	Health services research
Secondary Subject Heading:	Health policy
Keywords:	Health & safety < HEALTH SERVICES ADMINISTRATION & MANAGEMENT, Risk management < HEALTH SERVICES ADMINISTRATION & MANAGEMENT, Quality in health care < HEALTH SERVICES ADMINISTRATION & MANAGEMENT

SCHOLARONE™
Manuscripts

Unexpectedly long hospital stays as an indicator of risk of unsafe care: an exploratory study.

Ine Borghans ^{1,2§}, Karin D Hekkert ³, Lya den Ouden ², Sezgin Cihangir ³, Jan Vesseur ², Rudolf B Kool ¹, Gert P Westert ¹

¹ Scientific Institute for Quality of Healthcare (IQ healthcare), Radboud university medical centre, P.O. Box 9101, 6500 HB, Nijmegen, the Netherlands

- Ine Borghans, Senior research fellow
- Rudolf B Kool, Senior research fellow
- Gert P Westert, Head of the department

² Dutch Health Care Inspectorate (IGZ), P.O. Box 2680, 3500 GR, Utrecht, the Netherlands

- Ine Borghans, Coordinator Risk Analyzes
- Lya den Ouden, Coordinating specialistic Inspector
- Jan Vesseur, Project Chief Inspector Patient Safety

³ Dutch Hospital Data (DHD), P.O. Box 9696, 3506 GR Utrecht, the Netherlands

- Karin D Hekkert, Researcher
- Sezgin Cihangir, Researcher

[§] Corresponding author

Tijn Kool, P.O. Box 9101, 6500 HB, Nijmegen, the Netherlands
T: +31(0)611531143
E: tijn.kool@radboudumc.nl

Keywords: quality indicator, adverse events, length of stay, patient safety, hospital care

Word count : 3077

ABSTRACT

Objectives: We developed an outcome indicator based on the finding that complications often prolong the patient's hospital stay. A higher percentage of patients with an unexpectedly long length of stay (UL-LOS) compared to the national average may indicate shortcomings in patient safety. We explored the utility of the UL-LOS indicator.

Setting: We used data of 61 Dutch hospitals. In total these hospitals had 1,400,000 clinical discharges in 2011.

Participants: The indicator is based on the percentage of patients with a prolonged length of stay of more than 50% of the expected length of stay and calculated among survivors.

Interventions: no interventions were done.

Outcome measures: The outcome measures were the variability of the indicator across hospitals, the stability over time, the correlation between the UL-LOS and standardised mortality and the influence on the indicator of hospitals that did have problems discharging their patients to other health services such as nursing homes.

Results: In order to compare hospitals properly the expected length of stays was computed based on comparison with benchmark populations. The standardisation was based on patients' age, primary diagnosis and main procedure. The UL-LOS indicator showed considerable variability between the Dutch hospitals: from 8.6% to 20.1% in 2011. The outcomes had relatively small confidence intervals since they were based on large numbers of patients. The stability of the indicator over time was quite high. The indicator had a significant positive correlation with the standardised mortality ($r=0.44$ ($p<0.001$)), and no significant correlation with the percentage of patients that was discharged to other facilities than other hospitals and home ($r=-0.15$ ($p>0.05$)).

Conclusions: The UL-LOS indicator is a useful addition to other patient safety indicators by revealing variation between hospitals and areas of possible patient safety improvement.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

STRENGTHS AND LIMITATIONS OF THIS STUDY

- This study provides a hospital-wide indicator that can be used in addition to mortality and readmissions rates in order to identify potential safety risks
- The current indicator adjusts for differences in age, principal diagnosis and procedures. But there are probably more variables involved in a prolonged hospital stay.
- The indicator currently ‘counts’ all patients from whom the actual length of stay exceeds the expected duration by 50% or more. It needs to be studied whether this cut-off point can be set separately for each patient group.

For peer review only

BACKGROUND

For about ten years, improving quality of care based on outcome indicators is seen as an essential component in optimising safety in healthcare. In the Netherlands, like in many other countries, a large number of indicators have been developed and introduced to monitor the quality and safety of hospital care[1-3]. Many of these indicators concern a specific patient group. There are also some general quality indicators that concern the whole hospital. The most important ones are: unexpectedly long length of stay (UL-LOS); unplanned readmissions and higher than expected mortality (measured by the Hospital Standardised Mortality Rate, HSMR).

According to the Dutch Healthcare Inspectorate (IGZ) these are especially of interest for identifying general patient safety risks in hospital care. It is important to measure all three indicators since a degree of substitution -or competition- between them is possible. For example, if a hospital tries to discharge patients too quickly, this policy could result in a higher percentage of unplanned readmissions. The indicator for unplanned readmissions, already used in several countries such as the UK and USA[4-7], is not yet available in the Netherlands. The HSMR has already been available in Dutch hospitals since 2006[8].

The UL-LOS did not exist until some years ago the IGZ was looking for more general patient safety indicators. Research shows that hospital adverse events often result in a longer length of stay[9-22]. In several other studies on adverse events, a long length of stay was used as an important trigger for selecting medical records[23-25]. So, if complications often prolong the patient's hospital stay, could an outcome indicator be developed such as the percentage of patients with an unexpectedly long length of stay (UL-LOS) compared to the national average? Such an outcome may indicate shortcomings in the quality or safety of care delivered by the hospital.

As far back as 1999 Silber et al. had already published research about an indicator called 'conditional length of stay'. This was based on length of stay data and took into account the fact that patient stays tend to become prolonged after complications. They developed this indicator by testing if length of stay distributions display an 'extended' pattern of decreasing hazards after a transition point. This would suggest that 'the longer a patient has stayed in the hospital, the longer a patient will likely stay in the hospital'. Or, alternatively, there is the possibility that 'the longer a patient has stayed in the hospital, the faster a patient will likely be discharged from the hospital'[26].

Based on these former ideas, the IGZ decided to introduce such an indicator of prolonged length of stay. In the current paper we explore the utility of this indicator by measuring the variability of the indicator across hospitals and the stability over time. We also examine the correlation between the two existing indicators of the model: the UL-LOS and the HSMR, as they are both supposed to be an indicator of risk of unsafe care. The HSMR focuses by definition on adverse events leading to hospital mortality. The UL-LOS is calculated among survivors and involves a much wider range of adverse events. It includes all adverse events leading to a substantial prolonged length of stay. We expected a positive relationship between the two indicators, as reduced quality of care leads to more adverse events, and more adverse events lead to more patients with prolonged hospitalisation as well as to more deaths.

METHODS

Data

To calculate the UL-LOS we used data that were routinely registered for administrative purposes. Using an existing registration minimalizes extra registration burden. These databases can be used to predict risks with similar discrimination to clinical databases[27]. The National Medical Registration (LMR) already exists for 50 years in the Netherlands and contains data of hospital admissions including medical data such as diagnosis and surgical procedures as well as patient-specific data such as age and hospital stay[28]. For the UL-LOS we especially used the variable ‘expected length of stay’, which is generated by indirect standardisation based on the following three patient characteristics, which are the most import characteristics for standardisation of length of stay data[29]:

- Age: Divided into 5 categories: 0, 1-14, 15-44, 45-64, 65+ years;
- Primary diagnosis: This is the main diagnosis that led to the admission; it includes about 1,000 diagnoses classified by the ICD9 in three digits;
- Morbidity group: Morbidity groups mainly divide patients with or without procedures. For patients with procedures the morbidity groups are made by unique combinations of a diagnosis and one or more procedures. On average it includes five procedure groups. Procedures are classified by the Dutch Classification System of Procedures.

Together these three parameters produced $5 \times 1,000 \times 5 = 25,000$ cells. Every year the national mean length of stay of each cell is taken as the expected length of stay for the patients with characteristics that belong to the same cell.

Definition

We used the expected length of stay to define an outcome indicator that suited the purpose of identifying adverse events. We used former research to choose a single cut-off percentage prolonged length of stay across all diagnoses and case mixes to distinguish between 'normal' variation in length of stay and variation in hospital stay that might be caused by complications and other patient safety issues[30]. In literature we found the use of a threshold of 75%[31, 32], which in fact was arbitrary chosen. We formulated the indicator 'Unexpectedly long length of stay' as the percentage of clinically admitted patients with an actual hospital stay that was more than 50% longer than expected[2]. We excluded from this indicator patients who died in hospital. We excluded these patients because of the interrelationship between length of stay and mortality which makes it difficult to interpret length of stay data if non-survivors are included[33].

We used a threshold of 50% for two reasons:

1. We especially wanted to include patients that stayed longer because of complications and adverse events, and not patients that just stayed a little bit longer in hospital because of variations in efficiency, see figure 1. The percentage should not be too small.
2. We analysed for the Netherlands the percentages of patients with a longer than expected length of stay for ten different threshold groups between 0 and 100%. We found that the percentages after the threshold of 50% seem to dip beneath 2%. See annex 1.

In combination with our first experiences with case studies that showed that the number of adverse events clearly increased from a longer than expected length of stay of 50% or more[30], we decided to use the threshold of 50% in order not to exclude too many patients with adverse events. In fact the threshold depends to what specificity or what false positive rates you are willing to accept. With a threshold of 50% we expect to find a higher number of adverse events than with a threshold of 75%, but we expect the proportion of adverse events will be lower. A higher threshold would mean more efficiency in finding adverse events.

Sample

We selected hospitals for which the indicator could be calculated. Hospitals had to participate in the LMR not only by registration of the diagnoses of clinical patients but also the procedures. For our study this resulted in the use of data of the year 2011 of 61 out of all 90 Dutch hospitals:

- 6 hospitals were excluded because they did not participate at all

- 5 hospitals were excluded because they participated for less than 50% of the year. The group of 79 hospitals after this step consisted of 73 hospitals that participated the whole year, 4 hospitals participating for 50-60% of the year and 2 hospitals participating for 70-80% of the year. All these six hospitals participating for 50-80% coded at least six months sequent.
- 18 hospitals were excluded because they did not register the procedures in the LMR. Among these hospitals were general and tertiary teaching hospitals, and they did not differ in size or region from the hospitals that could be included in the study.

See the flowchart in figure 2.

To optimize the comparability of hospitals, we stratified the sample into three groups: 32 general hospitals, 21 tertiary teaching hospitals (TTHs) and 8 university medical centres (UMCs). Within these groups, size, degree of specialisation, financing system and complexity of patients are comparable[34-36].

Analyses

To explore the utility of the indicator, we measured the variability of the indicator across hospitals and the stability over time. To find out whether the indicator is stable over time, we determined the correlation between the percentages per hospital in ‘2008 and 2009’, ‘2009 and 2010’, ‘2010 and 2011’ and ‘2008 and 2011’.

In order to analyse whether the indicator could identify risks of unsafe care, we correlated the results of the UL-LOS with the HSMRs for the year 2011. Therefore we calculated the Pearson correlations and the 2-tailed Sig. between the unexpectedly long lengths of stay and HSMRs. The HSMR consists of the quotients of observed mortality and expected mortality in 50 diagnostic groups (CCS) in which 80% of all hospital mortality took place. The expected mortality was based on the following characteristics of the patients: age, sex, CCS-subgroup, co morbidity (Charlson index), urgency, social deprivation, source organisation type, month and year.

As with the UL-LOS, the HSMR could only be calculated for hospitals that participated in the LMR. Some additional criteria were used in order to optimize reliability of the standardisation. To be included in the HSMR, hospitals had to:

- Avoid the use of vague diagnostic codes (this had to be less than 2% of the admissions);
- Perform an adequate registration of the urgency of the admission (more than 30% of the admissions had to be marked as urgent);
- Perform an adequate registration of the comorbidity of patients (the mean number of secondary diagnosis per admission had to be more than 0,5).

In addition, the HSMR had to count for more than 70% of all deaths in hospital, and the hospital had to have more than 50 expected deaths per year. All these additional criteria resulted in 58 hospitals remaining for our correlation study between UL-LOS and HSMR for the year 2011, see the flowchart in figure 2.

We investigated whether the UL-LOS indicator (and the HSMR) might be influenced by hospitals that, for local resource reasons, do not have as much access to nursing home beds, home care support, palliative care or hospices, and thus keep more of these patients in the hospital rather than discharging them to stay or die elsewhere. We addressed this by two different analyses:

- 1) We calculated the percentage of patients that was discharged to other facilities than other hospitals and home. This percentage approximately gives the extent to which hospitals are able to discharge patients for long-term or palliative care.
- 2) We re-ran the analysis restricted to diagnostic categories where most patients are discharged home without a diagnosis for which one might expect a palliative approach. To select this group, we excluded all CCS groups[37] with carcinoma.

RESULTS

In Figure 3 the percentage of unexpectedly long length of stay is given for the 61 hospitals in our study. The figure also shows the 95% confidence limits. A distinction has been made between general hospitals, TTHs and UMCs. The national median of the percentage of clinical admissions with a UL-LOS was 11.3%. The UMCs had a relatively high score on this indicator compared with the TTHs and general hospitals.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For the UMCs the variation of the percentages was between 12.0% and 20.1%, with a median of 15.1%. The TTHs varied between 9.4% and 15.0%, with a median of 11.3%. For the general hospitals the variation was between 8.6% and 16.0%, with a median of 11.1%. With an independent samples t-test we found no significant difference between TTHs and general hospitals, $t(53) = 0.16$; $p = .88$.

To explore the stability of the indicator, we calculated the correlation of the indicator between two subsequent years. The R^2 between 2008 and 2009 was 0.89, between 2009 and 2010 it was 0.86 and between 2010 and 2011 it was 0.90. Figure 4 shows the correlation between 2008 and 2011. The R^2 between these years was 0.70.

Table 1 shows for the year 2011 for each hospital the UL-LOS indicator and the HSMR in two ways: with and without patients with carcinoma. This table also shows for each hospital the percentage of patients admitted to other destinations than home or other hospital. There was no substantial difference in the outcome of the UL-LOS indicator calculated with or without patients with carcinomas. The correlation of the UL-LOS with/without carcinomas was 1.00; correlation of the HSMR with/without carcinomas was 0.96. The Pearson correlation between the UL-LOS indicator and the HSMR with all diagnosis groups was 0.44 ($p < 0.001$) and with carcinoma excluded 0.52 ($p < 0.001$). We found no significant correlation between the UL-LOS indicator and the score on discharging patients. The Pearson correlation between the UL-LOS and discharging palliative patients with all diagnosis groups was -0.15 ($p > 0.05$) and with carcinoma excluded -0.16 ($p > 0.05$).

Table 1. 61 hospitals with their UL-LOS and HSMR in two ways: with and without patients with carcinoma and the percentage of patients admitted to other destinations than home or other hospital; 2011.

Hospital number	Total hospital		Carcinoma excluded		admissions to other destinations than home or other hospital
	UL-LOS 2011	HSMR 2011	UL-LOS 2011	HSMR 2011	
1	20,1%	117	20,3%	128	1,4%
2	18,7%	not available	18,8%	not available	0,6%
3	17,8%	115	17,8%	118	0,5%
4	16,0%	147	15,5%	145	3,4%
5	15,6%	118	15,6%	128	1,8%
6	15,6%	129	15,8%	133	1,4%
7	15,0%	100	15,0%	101	4,7%
8	14,7%	101	14,5%	103	0,9%
9	14,7%	not available	14,6%	not available	2,5%
10	14,7%	101	14,6%	100	4,3%
11	13,9%	93	13,6%	98	1,7%
12	13,6%	108	13,4%	105	3,2%
13	13,4%	113	13,3%	109	2,3%
14	13,4%	106	13,6%	104	not available
15	13,2%	71	13,0%	70	2,2%
16	13,0%	107	12,9%	112	1,2%
17	12,8%	112	12,6%	112	4,5%
18	12,7%	100	12,7%	101	1,7%
19	12,4%	108	12,2%	107	2,2%
20	12,4%	65	12,1%	64	3,1%
21	12,4%	106	12,2%	102	2,3%
22	12,3%	92	12,1%	88	2,1%
23	12,2%	70	12,1%	72	1,4%
24	12,1%	94	12,1%	86	2,1%
25	12,0%	91	12,3%	96	2,7%
26	11,7%	104	11,6%	102	0,2%
27	11,5%	78	11,5%	82	3,4%
28	11,5%	116	11,4%	111	1,2%
29	11,4%	115	11,5%	114	2,6%
30	11,4%	101	11,1%	105	1,0%
31	11,3%	101	11,0%	90	0,8%
32	11,3%	94	11,3%	96	2,8%
33	11,3%	94	11,3%	98	2,8%
34	11,3%	107	11,3%	115	2,3%
35	11,2%	91	11,2%	91	3,3%

36	11,2%	83	11,1%	80	0,6%
37	11,1%	104	11,3%	106	1,5%
38	11,1%	114	10,8%	111	1,8%
39	11,0%	103	10,9%	108	2,4%
40	11,0%	93	10,7%	91	2,2%
41	10,9%	114	10,9%	114	2,4%
42	10,9%	106	10,8%	103	2,9%
43	10,9%	107	10,8%	105	2,0%
44	10,9%	104	10,5%	101	2,5%
45	10,7%	110	10,6%	101	3,4%
46	10,7%	81	10,6%	79	2,4%
47	10,6%	100	10,4%	99	1,6%
48	10,6%	99	10,6%	95	2,4%
49	10,6%	not available	10,2%	not available	2,2%
50	10,5%	93	10,3%	95	2,3%
51	10,4%	91	10,1%	88	3,1%
52	10,1%	107	10,0%	103	3,0%
53	10,1%	81	10,4%	78	2,0%
54	9,9%	101	9,7%	104	3,1%
55	9,9%	61	9,6%	65	3,8%
56	9,7%	111	9,7%	112	1,2%
57	9,6%	99	9,5%	101	2,9%
58	9,4%	82	9,5%	85	3,1%
59	9,4%	83	9,2%	77	1,3%
60	9,3%	89	9,3%	86	2,7%
61	8,6%	91	8,7%	93	1,0%

DISCUSSION

In this paper, we described the development of a patient safety indicator for Dutch hospitals that, to the best of our knowledge, has not been described in the literature till now. The indicator is defined as the percentage of clinically admitted patients with an actual hospital stay that was more than 50% longer than expected. The indicator showed considerable variability between the Dutch hospitals: from 8.6% to 20.1% in 2011. It also showed serious variation within homogenous groups of hospitals. The stability of the indicator over three years was quite high and the indicator had a significant positive correlation with the HSMR. This indicates that the hospitals with more patients with a UL-LOS were also the hospitals with higher standardised mortality. This might support our hypothesis that sub-optimal quality of care may both lead to more patients with an unexpectedly long length of stay as well as to more mortality than expected. More research is needed to determine the

validity of the UL-LOS indicator. Especially record reviewing to look for adverse events is needed for a stronger support of our hypothesis. In addition to this an indicator for readmissions should be added to the indicator framework and should be subject for future research.

The strong correlation between the UL-LOS with and without carcinomas, and the low correlation between UL-LOS and discharging patients, seem to indicate that the UL-LOS is not influenced by differences between hospitals in the ability to admit patients to next care facilities. This might be important for hospitals in realizing that high percentages of patients with an UL-LOS will probably not be caused by opportunities of discharging patients.

In terms of evaluation of care it is becoming increasingly common for hospitals to study patient records retrospectively, especially in cases of deceased patients^{25,26}. This indicator might be a good research tool to identify records of patients who were discharged alive. This could lead to opportunities for improvements that are different from those based on patient record reviewing after death.

The indicator provides insight into the percentage of patients that stayed at least 50% longer than expected. The assumption is that in this group a relatively large number of patients have had to deal with unexpected developments in their disease, resulting in complications that cause a prolonged stay. It could be much more effective for hospitals to review records of hospital admissions selected by this indicator compared to randomly selected patient records. Reviewing records takes considerable time and by using this indicator for selection, time could be saved by reviewing fewer records from which more lessons might be learnt[30].

Hospital management might also have financial reasons to be interested in the indicator in addition to the quality and safety aspects of the UL-LOS indicator. Having a longer than expected length of stay, costs money in the present Dutch financing system. Although the indicator is not developed for financial purposes, the use of the indicator may, as a beneficial side effect, also reduce the total amount of hospital days.

There are two main limitations to this study.

1. Some hospitals, especially the UMCs, differed from the national median. The high score for the UMCs could indicate that there is insufficient adjustment for the specific patient categories admitted to the UMCs. The current indicator adjusts for differences in age, principal diagnosis and procedures. But there are probably more variables involved in a prolonged hospital stay. The case mix

adjustment is more limited than for example the HSMR. Further research is needed to determine which other patient characteristics, e.g. co-morbidity, play a significant role in a prolonged length of stay and whether they are of added value in standardising hospital stay next to the present co-founders.

2. We used a threshold of a length of stay of 50% longer than expected for the indicator. This threshold was based on the three above mentioned reasons and especially our first experiences with reviewing hospital records based on length of stay[30]. However, there is no evidence that the threshold has to be exactly 50%. A more detailed study is needed to determine the appropriateness of this threshold. Further research is also needed to determine to what extent the proportion of patients which crosses the threshold can vary for each combination of age, main diagnosis and procedure. If the variations between hospitals are large, and the case mix clearly differs, this will present a number of unequal opportunities that will result in crossing the 50% threshold. We might need to vary the threshold for different diagnostic groups.

CONCLUSION

The IGZ introduced the outcome indicator 'Percentage of patients with an unexpectedly long length of stay' for its supervision since 2010. It is based on the assumption that complications often prolong the patient's hospital stay. A higher percentage of patients with a UL-LOS compared to the national average, after a correction is made for case mix variations, may indicate shortcomings in the quality or safety of care delivered by the hospital. The indicator currently 'counts' all patients from whom the actual length of stay exceeds the expected duration by 50% or more. It needs to be studied whether this cut-off point can be set separately for each patient group. The indicator varies systematically between hospitals and is rather stable over time. It correlates with other outcome indicators, which could indicate the capacity to identify opportunities for improvement of patient safety.

FIGURE LEGENDS

Figure 1: Components of length of stay

Figure 1: LOS = length of stay
 UL-LOS = unexpectedly long length of stay

Figure 2: Flowchart hospitals in correlation study between UL-LOS and HSMR for the year 2011

Figure 2: LMR = National Medical Registration
 HSMR = Hospital Standardised Mortality Rate

Figure 3. Percentage of admissions with an unexpectedly long length of stay (UL-LOS) for the hospitals in our study, defined by type of hospital: general hospitals (n=32), tertiary teaching hospitals (n=21) and University Medical Centres (n=8); LMR 2011

Figure 3: I = 95% CI
 • = Admissions with UL-LOS 2011 (%)

Figure 4. Correlation between the years 2008 and 2011 for the percentage of admissions with a UL-LOS per hospital. LMR 2008 and 2011 (n=57)

Figure 4: • = hospital
 UL-LOS = unexpectedly long length of stay

ACKNOWLEDGMENTS

COMPETING INTERESTS

The authors declare that they have no competing interests.

FUNDING

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

CONTRIBUTORSHIP STATEMENT

IB designed the study, performed the analysis, interpreted the results and drafted the manuscript. KDH and SC contributed to the analysis and the interpretation of the results. RBK, LdO and JV helped to interpret the results and contributed to the discussion. RBK and GPW supervised the study and participated in the formulation of the discussion. All authors reviewed and edited the manuscript for intellectual content. All authors read and approved the final manuscript.

DATA SHARING STATEMENT

No additional data are available

REFERENCES

1 Inspectie voor de Gezondheidszorg [Dutch Health Care Inspectorate]. *Basisset kwaliteitsindicatoren ziekenhuizen 2013*. Utrecht: IGZ; 2012.

2 Inspectie voor de Gezondheidszorg [Dutch Health Care Inspectorate]. *Basisset veiligheidsindicatoren ziekenhuizen 2010-2012*. Utrecht: IGZ; 2010.

3 Kwaliteitsindicatoren Zichtbare Zorg Ziekenhuizen. Available at: <http://www.zichtbarezorg.nl/page/Ziekenhuizen-en-ZBC-s/Kwaliteitsindicatoren#ZI>. Accessed 12/01, 2011.

4 Nolte E, Roland M, Guthrie S, et al. Preventing emergency readmissions to hospital, A scoping review. RAND Europe; 2012.

5 Lagoe RJ, Nanno DS, Luziani ME. Quantitative tools for addressing hospital readmissions. *BMC Res Notes* 2012;**5**:620,0500-5-620.

6 Goldfield NI, McCullough EC, Hughes JS, et al. Identifying potentially preventable readmissions. *Health Care Financ Rev* 2008;**30**:75-91.

7 Joynt KE, Jha AK. Thirty-day readmissions--truth and consequences. *N Engl J Med* 2012;**366**:1366-9.

8 Jarman B, Pieter D, van der Veen A, et al. The hospital standardised mortality ratio: a powerful tool for Dutch hospitals to assess their quality of care?. *Qual Saf Health Care* 2010;**19**:9-13.

9 Hoonhout LH, de Bruijne MC, Wagner C, et al. Nature, occurrence and consequences of medication-related adverse events during hospitalization: a retrospective chart review in the Netherlands. *Drug Saf* 2010;**33**:853-64.

10 Hoonhout LH, de Bruijne MC, Wagner C, et al. Direct medical costs of adverse events in Dutch hospitals. *BMC Health Serv Res* 2009;**9**:27.

11 Sari AB, Sheldon TA, Cracknell A, et al. Extent, nature and consequences of adverse events: results of a retrospective casenote review in a large NHS hospital. *Qual Saf Health Care* 2007;**16**:434-9.

12 Ehsani JP, Jackson T, Duckett SJ. The incidence and cost of adverse events in Victorian hospitals 2003-04. *Med J Aust* 2006;**184**:551-5.

13 Cho SH, Ketefian S, Barkauskas VH, et al. The effects of nurse staffing on adverse events, morbidity, mortality, and medical costs. *Nurs Res* 2003;**52**:71-9.

14 Camp M, Chang DC, Zhang Y, et al. Risk factors and outcomes for foreign body left during a procedure: analysis of 413 incidents after 1 946 831 operations in children. *Arch Surg* 2010;**145**:1085-90.

15 Lotfipour S, Kaku SK, Vaca FE, et al. Factors associated with complications in older adults with isolated blunt chest trauma. *West J Emerg Med* 2009;**10**:79-84.

-
- 16 Williams DJ, Olsen S, Crichton W, et al. Detection of adverse events in a Scottish hospital using a consensus-based methodology. *Scott Med J* 2008;**53**:26-30.
- 17 Kaushal R, Bates DW, Franz C, et al. Costs of adverse events in intensive care units. *Crit Care Med* 2007;**35**:2479-83.
- 18 Rice-Townsend S, Hall M, Jenkins KJ, et al. Analysis of adverse events in pediatric surgery using criteria validated from the adult population: justifying the need for pediatric-focused outcome measures. *J Pediatr Surg* 2010;**45**:1126-36.
- 19 Schioler T, Lipczak H, Pedersen BL, et al. Incidence of adverse events in hospitals. A retrospective study of medical records. *Ugeskr Laeger* 2001;**163**:5370-8.
- 20 Baker GR, Norton PG, Flintoft V, et al. The Canadian Adverse Events Study: the incidence of adverse events among hospital patients in Canada. *CMAJ* 2004;**170**:1678-86.
- 21 Davis P, Lay-Yee R, Briant R, et al. Adverse events in New Zealand public hospitals I: occurrence and impact. *N Z Med J* 2002;**115**:U271.
- 22 Lagoe R, Pernisi L, Luziani M, et al. Addressing hospital length of stay outlier patients: A community wide approach. *Advances in Bioscience and Biotechnology* 2014;**6**:188-96.
- 23 de Bruijne MC, Zegers M, Hoonhout LHF, et al. Onbedoelde schade in Nederlandse ziekenhuizen: dossieronderzoek van ziekenhuisopnames in 2004. Amsterdam: EMGO Instituut/VUmc en Nivel, Nederlands Instituut voor onderzoek van de gezondheidszorg; 2007.
- 24 Langelan M, Baines RJ, Broekens MA, et al. Monitor Zorggerelateerde Schade 2008, dossieronderzoek in Nederlandse ziekenhuizen. NIVEL; 2010.
- 25 Zegers M, de Bruijne MC, Wagner C, et al. Adverse events and potentially preventable deaths in Dutch hospitals: results of a retrospective patient record review study. *Qual Saf Health Care* 2009;**18**:297-302.

26 Silber JH, Rosenbaum PR, Koziol LF, et al. Conditional Length of Stay. *Health Serv Res* 1999;**34**:349-63.

27 Aylin P, Bottle A, Majeed A. Use of administrative data or clinical databases as predictors of risk of death in hospital: comparison of models. *BMJ* 2007;**334**:1044.

28 Landelijke Medische Registratie (LMR). Available at:
<http://www.dutchhospitaldata.nl/registraties/lmrlazr/Paginas/default.aspx>. Accessed 06/10, 2013.

29 Hart 't P, de Boer K, Draaisma D. On the length of stay in hospitals. Radboud University Nijmegen; 2012.

30 Cihangir S, Borghans I, Hekkert K, et al. A pilot study on record reviewing with a priori patient selection. *BMJ Open* 2013;**3**:10.1136/bmjopen.2013-003034. Print 2013.

31 Bottle A, Aylin P. Intelligent information: a national system for monitoring clinical performance. *Health Serv Res* 2008;**43**:10-31.

32 Bottle A, Middleton S, Kalkman CJ, et al. Global comparators project: international comparison of hospital outcomes using administrative data. *Health Serv Res* 2013;**48**:2081-100.

33 Marang-van de Mheen PJ, Lingsma HF, Middleton S, et al. Evaluation of quality of care using registry data: the interrelationship between length-of-stay, readmission and mortality and impact on hospital outcomes. *BMJ Qual Saf* 2014;**23**:350-1.

34 Website of more than 40 small and medium-sized Dutch hospitals. They take more than a third of all Dutch hospital care on their behalf. Available at: <http://www.saz-ziekenhuizen.nl>. Accessed 09-30, 2013.

35 Website with facts and figures of all (8) Dutch university hospitals. Available at:
<http://www.nfu.nl/umc/feiten/>. Accessed 09-30, 2013.

36 Website of 28 Dutch tertiary teaching hospitals. Available at: <http://www.stz.nl/pagina/9-over-stz.html>.
Accessed 09-30, 2013.

37 Rivard PE, Luther SL, Christiansen CL, et al. Using patient safety indicators to estimate the impact of
potential adverse events on outcomes. *Med Care Res Rev* 2008;**65**:67-87.

**Unexpectedly long hospital stays as an indicator of risk of unsafe care:
an exploratory study.**

Ine Borghans ^{1,2§}, Karin D Hekkert ³, Lya den Ouden ², Sezgin Cihangir ³, Jan Vesseur ²,
Rudolf B Kool ¹, Gert P Westert ¹

¹ Scientific Institute for Quality of Healthcare (IQ healthcare), Radboud university
medical centre, P.O. Box 9101, 6500 HB, Nijmegen, the Netherlands

- Ine Borghans, Senior research fellow
- Rudolf B Kool, Senior research fellow
- Gert P Westert, Head of the department

² Dutch Health Care Inspectorate (IGZ), P.O. Box 2680, 3500 GR, Utrecht, the
Netherlands

- Ine Borghans, Coordinator Risk Analyzes
- Lya den Ouden, Coordinating specialistic Inspector
- Jan Vesseur, Project Chief Inspector Patient Safety

³ Dutch Hospital Data (DHD), P.O. Box 9696, 3506 GR Utrecht, the Netherlands

- Karin D Hekkert, Researcher
- Sezgin Cihangir, Researcher

[§] Corresponding author

Tijn Kool, P.O. Box 9101, 6500 HB, Nijmegen, the Netherlands

T: +31(0)611531143

E: tijn.kool@radboudumc.nl

Keywords: quality indicator, adverse events, length of stay, patient safety, hospital
care

Word count : 3077

ABSTRACT

Objectives: We developed an outcome indicator based on the finding that complications often prolong the patient's hospital stay. A higher percentage of patients with an unexpectedly long length of stay (UL-LOS) compared to the national average may indicate shortcomings in patient safety. We explored the utility of the UL-LOS indicator.

Setting: We used data of 61 Dutch hospitals. In total these hospitals had 1,400,000 clinical discharges in 2011.

Participants: The indicator is based on the percentage of patients with a prolonged length of stay of more than 50% of the expected length of stay and calculated among survivors.

Interventions: no interventions were done.

Outcome measures: The outcome measures were the variability of the indicator across hospitals, the stability over time, the correlation between the UL-LOS and standardised mortality and the influence on the indicator of hospitals that did have problems discharging their patients to other health services such as nursing homes.

Results: In order to compare hospitals properly the expected length of stays was computed based on comparison with benchmark populations. The standardisation was based on patients' age, primary diagnosis and main procedure. The UL-LOS indicator showed considerable variability between the Dutch hospitals: from 8.6% to 20.1% in 2011. The outcomes had relatively small confidence intervals since they were based on large numbers of patients. The stability of the indicator over time was quite high. The indicator had a significant positive correlation with the standardised mortality ($r=0.44$ ($p<0.001$)), and no significant correlation with the percentage of patients that was discharged to other facilities than other hospitals and home ($r=-0.15$ ($p>0.05$)).

Conclusions: The UL-LOS indicator is a useful addition to other patient safety indicators by revealing variation between hospitals and areas of possible patient safety improvement.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

STRENGTHS AND LIMITATIONS OF THIS STUDY

- This study provides a hospital-wide indicator that can be used in addition to mortality and readmissions rates in order to identify potential safety risks
- The current indicator adjusts for differences in age, principal diagnosis and procedures. But there are probably more variables involved in a prolonged hospital stay.
- The indicator currently ‘counts’ all patients from whom the actual length of stay exceeds the expected duration by 50% or more. It needs to be studied whether this cut-off point can be set separately for each patient group.

For peer review only

BACKGROUND

For about ten years, improving quality of care based on outcome indicators is seen as an essential component in optimising safety in healthcare. In the Netherlands, like in many other countries, a large number of indicators have been developed and introduced to monitor the quality and safety of hospital care[1-3]. Many of these indicators concern a specific patient group. There are also some general quality indicators that concern the whole hospital. The most important ones are: unexpectedly long length of stay (UL-LOS); unplanned readmissions and higher than expected mortality (measured by the Hospital Standardised Mortality Rate, HSMR).

According to the Dutch Healthcare Inspectorate (IGZ) these are especially of interest for identifying general patient safety risks in hospital care. It is important to measure all three indicators since a degree of substitution -or competition- between them is possible. For example, if a hospital tries to discharge patients too quickly, this policy could result in a higher percentage of unplanned readmissions. The indicator for unplanned readmissions, already used in several countries such as the UK and USA[4-7], is not yet available in the Netherlands. The HSMR has already been available in Dutch hospitals since 2006[8].

The UL-LOS did not exist until some years ago the IGZ was looking for more general patient safety indicators. Research shows that hospital adverse events often result in a longer length of stay[9-22]. In several other studies on adverse events, a long length of stay was used as an important trigger for selecting medical records[23-25]. So, if complications often prolong the patient's hospital stay, could an outcome indicator be developed such as the percentage of patients with an unexpectedly long length of stay (UL-LOS) compared to the national average? Such an outcome may indicate shortcomings in the quality or safety of care delivered by the hospital.

As far back as 1999 Silber et al. had already published research about an indicator called 'conditional length of stay'. This was based on length of stay data and took into account the fact that patient stays tend to become prolonged after complications. They developed this indicator by testing if length of stay distributions display an 'extended' pattern of decreasing hazards after a transition point. This would suggest that 'the longer a patient has stayed in the hospital, the longer a patient will likely stay in the hospital'. Or, alternatively, there is the possibility that 'the longer a patient has stayed in the hospital, the faster a patient will likely be discharged from the hospital'[26].

Based on these former ideas, the IGZ decided to introduce such an indicator of prolonged length of stay. In the current paper we explore the utility of this indicator by measuring the variability of the indicator across hospitals and the stability over time. We also examine the correlation between the two existing indicators of the model: the UL-LOS and the HSMR, as they are both supposed to be an indicator of risk of unsafe care. The HSMR focuses by definition on adverse events leading to hospital mortality. The UL-LOS is calculated among survivors and involves a much wider range of adverse events. It includes all adverse events leading to a substantial prolonged length of stay. We expected a positive relationship between the two indicators, as reduced quality of care leads to more adverse events, and more adverse events lead to more patients with prolonged hospitalisation as well as to more deaths.

METHODS

Data

To calculate the UL-LOS we used data that were routinely registered for administrative purposes. Using an existing registration minimalizes extra registration burden. These databases can be used to predict risks with similar discrimination to clinical databases[27]. The National Medical Registration (LMR) already exists for 50 years in the Netherlands and contains data of hospital admissions including medical data such as diagnosis and surgical procedures as well as patient-specific data such as age and hospital stay[28]. For the UL-LOS we especially used the variable ‘expected length of stay’, which is generated by indirect standardisation based on the following three patient characteristics, which are the most import characteristics for standardisation of length of stay data[29]:

- Age: Divided into 5 categories: 0, 1-14, 15-44, 45-64, 65+ years;
- Primary diagnosis: This is the main diagnosis that led to the admission; it includes about 1,000 diagnoses classified by the ICD9 in three digits;
- Morbidity group: Morbidity groups mainly divide patients with or without procedures. For patients with procedures the morbidity groups are made by unique combinations of a diagnosis and one or more procedures. On average it includes five procedure groups. Procedures are classified by the Dutch Classification System of Procedures.

Together these three parameters produced $5 \times 1,000 \times 5 = 25,000$ cells. Every year the national mean length of stay of each cell is taken as the expected length of stay for the patients with characteristics that belong to the same cell.


Definition

We used the expected length of stay to define an outcome indicator that suited the purpose of identifying adverse events. We used former research to choose a single cut-off percentage prolonged length of stay across all diagnoses and case mixes to distinguish between 'normal' variation in length of stay and variation in hospital stay that might be caused by complications and other patient safety issues[30]. In literature we found the use of a threshold of 75%[31, 32], which in fact was arbitrary chosen. We formulated the indicator 'Unexpectedly long length of stay' as the percentage of clinically admitted patients with an actual hospital stay that was more than 50% longer than expected[2]. We excluded from this indicator patients who died in hospital. We excluded these patients because of the interrelationship between length of stay and mortality which makes it difficult to interpret length of stay data if non-survivors are included[33].

We used a threshold of 50% for two reasons:

1. We especially wanted to include patients that stayed longer because of complications and adverse events, and not patients that just stayed a little bit longer in hospital because of variations in efficiency, see figure 1. The percentage should not be too small.

Figure 1: Components of length of stay



2. We analysed for the Netherlands the percentages of patients with a longer than expected length of stay for ten different threshold groups between 0 and 100%. We found that the percentages after the threshold of 50% seem to dip beneath 2%. See annex 1.

In combination with our first experiences with case studies that showed that the number of adverse events clearly increased from a longer than expected length of stay of 50% or more[30], we decided to use the threshold of 50% in order not to exclude too many patients with adverse events. In fact the threshold depends to what specificity or what false positive rates you are willing to accept. With a threshold of 50% we expect to find a higher number of adverse events than with a threshold of 75%, but we expect the proportion of adverse events will be lower. A higher threshold would mean more efficiency in finding adverse events.

Sample

We selected hospitals for which the indicator could be calculated. Hospitals had to participate in the LMR not only by registration of the diagnoses of clinical patients but also the procedures. For our study this resulted in the use of data of the year 2011 of 61 out of all 90 Dutch hospitals:

- 6 hospitals were excluded because they did not participate at all
- 5 hospitals were excluded because they participated for less than 50% of the year. The group of 79 hospitals after this step consisted of 73 hospitals that participated the whole year, 4 hospitals participating for 50-60% of the year en 2 hospitals participating for 70-80% of the year. All these six hospitals participating for 50-80% coded at least six months sequent.
- 18 hospitals were excluded because they did not register the procedures in the LMR. Among these hospitals were general and tertiary teaching hospitals, and they did not differ in size or region from the hospitals that could be included in the study.

See the flowchart in figure 2.

To optimize the comparability of hospitals, we stratified the sample into three groups: 32 general hospitals, 21 tertiary teaching hospitals (TTHs) and 8 university medical centres (UMCs). Within these groups, size, degree of specialisation, financing system and complexity of patients are comparable[34-36].

Analyses

To explore the utility of the indicator, we measured the variability of the indicator across hospitals and the stability over time. To find out whether the indicator is stable over time, we determined the correlation between the percentages per hospital in ‘2008 and 2009’, ‘2009 and 2010’, ‘2010 and 2011’ and ‘2008 and 2011’.

In order to analyse whether the indicator could identify risks of unsafe care, we correlated the results of the UL-LOS with the HSMRs for the year 2011. Therefore we


calculated the Pearson correlations and the 2-tailed Sig. between the unexpectedly long lengths of stay and HSMRs. The HSMR consists of the quotients of observed mortality and expected mortality in 50 diagnostic groups (CCS) in which 80% of all hospital mortality took place. The expected mortality was based on the following characteristics of the patients: age, sex, CCS-subgroup, co morbidity (Charlson index), urgency, social deprivation, source organisation type, month and year.

As with the UL-LOS, the HSMR could only be calculated for hospitals that participated in the LMR. Some additional criteria were used in order to optimize reliability of the standardisation. To be included in the HSMR, hospitals had to:

- Avoid the use of vague diagnostic codes (this had to be less than 2% of the admissions);
- Perform an adequate registration of the urgency of the admission (more than 30% of the admissions had to be marked as urgent);
- Perform an adequate registration of the comorbidity of patients (the mean number of secondary diagnosis per admission had to be more than 0,5).

In addition, the HSMR had to count for more than 70% of all deaths in hospital, and the hospital had to have more than 50 expected deaths per year. All these additional criteria resulted in 58 hospitals remaining for our correlation study between UL-LOS and HSMR for the year 2011, [see the flowchart in figure 2](#).

Figure 2: Flowchart hospitals in correlation study between UL-LOS and HSMR for the year 2011




We investigated whether the UL-LOS indicator (and the HSMR) might be influenced by hospitals that, for local resource reasons, do not have as much access to nursing home beds, home care support, palliative care or hospices, and thus keep more of these patients in the hospital rather than discharging them to stay or die elsewhere. We addressed this by two different analyses:

- 1) We calculated the percentage of patients that was discharged to other facilities than other hospitals and home. This percentage approximately gives the extent to which hospitals are able to discharge patients for long-term or palliative care.
- 2) We re-ran the analysis restricted to diagnostic categories where most patients are discharged home without a diagnosis for which one might expect a palliative approach. To select this group, we excluded all CCS groups[37] with carcinoma.

RESULTS

In Figure 3 the percentage of unexpectedly long length of stay is given for the 61 hospitals in our study. The figure also shows the 95% confidence limits. A distinction has been made between general hospitals, TTHs and UMCs. The national median of the percentage of clinical admissions with a UL-LOS was 11.3%. The UMCs had a relatively high score on this indicator compared with the TTHs and general hospitals.

Figure 3. Percentage of admissions with an unexpectedly long length of stay (UL-LOS) for the hospitals in our study, defined by type of hospital: general hospitals (n=32), tertiary teaching hospitals (n=21) and University Medical Centres (n=8); LMR 2011



For the UMCs the variation of the percentages was between 12.0% and 20.1%, with a median of 15.1%. The TTHs varied between 9.4% and 15.0%, with a median of 11.3%. For the general hospitals the variation was between 8.6% and 16.0%, with a median of 11.1%. With an independent samples t-test we found no significant difference between TTHs and general hospitals, $t(53) = 0.16$; $p = .88$.

To explore the stability of the indicator, we calculated the correlation of the indicator between two subsequent years. The R^2 between 2008 and 2009 was 0.89, between 2009 and 2010 it was 0.86 and between 2010 and 2011 it was 0.90. Figure 4 shows the correlation between 2008 and 2011. The R^2 between these years was 0.70.

Figure 4. Correlation between the years 2008 and 2011 for the percentage of admissions with a UL-LOS per hospital. LMR 2008 and 2011 (n=57)




Table 1 shows for the year 2011 for each hospital the UL-LOS indicator and the HSMR in two ways: with and without patients with carcinoma. This table also shows for each hospital the percentage of patients admitted to other destinations than home or other hospital. There was no substantial difference in the outcome of the UL-LOS indicator calculated with or without patients with carcinomas. The correlation of the UL-LOS with/without carcinomas was 1.00; correlation of the HSMR with/without carcinomas was 0.96. The Pearson correlation between the UL-LOS indicator and the HSMR with all diagnosis groups was 0.44 ($p < 0.001$) and with carcinoma excluded 0.52 ($p < 0.001$). We found no significant correlation between the UL-LOS indicator and the score on discharging patients. The Pearson correlation between the UL-LOS and discharging palliative patients with all diagnosis groups was -0.15 ($p > 0.05$) and with carcinoma excluded -0.16 ($p > 0.05$).

Table 1. 61 hospitals with their UL-LOS and HSMR in two ways: with and without patients with carcinoma and the percentage of patients admitted to other destinations than home or other hospital; 2011.

	Total hospital		Carcinoma excluded		
Hospital number	UL-LOS 2011	HSMR 2011	UL-LOS 2011	HSMR 2011	admissions to other destinations than home or other hospital
1	20,1%	117	20,3%	128	1,4%
2	18,7%	not available	18,8%	not available	0,6%
3	17,8%	115	17,8%	118	0,5%
4	16,0%	147	15,5%	145	3,4%
5	15,6%	118	15,6%	128	1,8%
6	15,6%	129	15,8%	133	1,4%
7	15,0%	100	15,0%	101	4,7%
8	14,7%	101	14,5%	103	0,9%
9	14,7%	not available	14,6%	not available	2,5%
10	14,7%	101	14,6%	100	4,3%
11	13,9%	93	13,6%	98	1,7%
12	13,6%	108	13,4%	105	3,2%
13	13,4%	113	13,3%	109	2,3%
14	13,4%	106	13,6%	104	not available
15	13,2%	71	13,0%	70	2,2%
16	13,0%	107	12,9%	112	1,2%
17	12,8%	112	12,6%	112	4,5%
18	12,7%	100	12,7%	101	1,7%
19	12,4%	108	12,2%	107	2,2%
20	12,4%	65	12,1%	64	3,1%
21	12,4%	106	12,2%	102	2,3%
22	12,3%	92	12,1%	88	2,1%
23	12,2%	70	12,1%	72	1,4%
24	12,1%	94	12,1%	86	2,1%
25	12,0%	91	12,3%	96	2,7%
26	11,7%	104	11,6%	102	0,2%
27	11,5%	78	11,5%	82	3,4%
28	11,5%	116	11,4%	111	1,2%
29	11,4%	115	11,5%	114	2,6%
30	11,4%	101	11,1%	105	1,0%
31	11,3%	101	11,0%	90	0,8%
32	11,3%	94	11,3%	96	2,8%
33	11,3%	94	11,3%	98	2,8%
34	11,3%	107	11,3%	115	2,3%
35	11,2%	91	11,2%	91	3,3%

36	11,2%	83	11,1%	80	0,6%
37	11,1%	104	11,3%	106	1,5%
38	11,1%	114	10,8%	111	1,8%
39	11,0%	103	10,9%	108	2,4%
40	11,0%	93	10,7%	91	2,2%
41	10,9%	114	10,9%	114	2,4%
42	10,9%	106	10,8%	103	2,9%
43	10,9%	107	10,8%	105	2,0%
44	10,9%	104	10,5%	101	2,5%
45	10,7%	110	10,6%	101	3,4%
46	10,7%	81	10,6%	79	2,4%
47	10,6%	100	10,4%	99	1,6%
48	10,6%	99	10,6%	95	2,4%
49	10,6%	not available	10,2%	not available	2,2%
50	10,5%	93	10,3%	95	2,3%
51	10,4%	91	10,1%	88	3,1%
52	10,1%	107	10,0%	103	3,0%
53	10,1%	81	10,4%	78	2,0%
54	9,9%	101	9,7%	104	3,1%
55	9,9%	61	9,6%	65	3,8%
56	9,7%	111	9,7%	112	1,2%
57	9,6%	99	9,5%	101	2,9%
58	9,4%	82	9,5%	85	3,1%
59	9,4%	83	9,2%	77	1,3%
60	9,3%	89	9,3%	86	2,7%
61	8,6%	91	8,7%	93	1,0%

DISCUSSION

In this paper, we described the development of a patient safety indicator for Dutch hospitals that, to the best of our knowledge, has not been described in the literature till now. The indicator is defined as the percentage of clinically admitted patients with an actual hospital stay that was more than 50% longer than expected. The indicator showed considerable variability between the Dutch hospitals: from 8.6% to 20.1% in 2011. It also showed serious variation within homogenous groups of hospitals. The stability of the indicator over three years was quite high and the indicator had a significant positive correlation with the HSMR. This indicates that the hospitals with more patients with a UL-LOS were also the hospitals with higher standardised mortality. This might support our hypothesis that sub-optimal quality of care may both lead to more patients with an unexpectedly long length of stay as well as to more mortality than expected. **More research is needed to determine the**

validity of the UL-LOS indicator. Especially record reviewing to look for adverse events is needed for a stronger support of our hypothesis. In addition to this an indicator for readmissions should be added to the indicator framework and should be subject for future research.

The strong correlation between the UL-LOS with and without carcinomas, and the low correlation between UL-LOS and discharging patients, seem to indicate that the UL-LOS is not influenced by differences between hospitals in the ability to admit patients to next care facilities. This might be important for hospitals in realizing that high percentages of patients with an UL-LOS will probably not be caused by opportunities of discharging patients.

In terms of evaluation of care it is becoming increasingly common for hospitals to study patient records retrospectively, especially in cases of deceased patients^{25,26}. This indicator might be a good research tool to identify records of patients who were discharged alive. This could lead to opportunities for improvements that are different from those based on patient record reviewing after death.

The indicator provides insight into the percentage of patients that stayed at least 50% longer than expected. The assumption is that in this group a relatively large number of patients have had to deal with unexpected developments in their disease, resulting in complications that cause a prolonged stay. It could be much more effective for hospitals to review records of hospital admissions selected by this indicator compared to randomly selected patient records. Reviewing records takes considerable time and by using this indicator for selection, time could be saved by reviewing fewer records from which more lessons might be learnt[30].

Hospital management might also have financial reasons to be interested in the indicator in addition to the quality and safety aspects of the UL-LOS indicator. Having a longer than expected length of stay, costs money in the present Dutch financing system. Although the indicator is not developed for financial purposes, the use of the indicator may, as a beneficial side effect, also reduce the total amount of hospital days.

There are two main limitations to this study.

1. Some hospitals, especially the UMCs, differed from the national median. The high score for the UMCs could indicate that there is insufficient adjustment for the specific patient categories admitted to the UMCs. The current indicator adjusts for differences in age, principal diagnosis and procedures. But there are probably more variables involved in a prolonged hospital stay. The case mix

adjustment is more limited than for example the HSMR. Further research is needed to determine which other patient characteristics, e.g. co-morbidity, play a significant role in a prolonged length of stay and whether they are of added value in standardising hospital stay next to the present co-founders.

2. We used a threshold of a length of stay of 50% longer than expected for the indicator. This threshold was based on the three above mentioned reasons and especially our first experiences with reviewing hospital records based on length of stay[30]. However, there is no evidence that the threshold has to be exactly 50%. A more detailed study is needed to determine the appropriateness of this threshold. Further research is also needed to determine to what extent the proportion of patients which crosses the threshold can vary for each combination of age, main diagnosis and procedure. If the variations between hospitals are large, and the case mix clearly differs, this will present a number of unequal opportunities that will result in crossing the 50% threshold. We might need to vary the threshold for different diagnostic groups.

CONCLUSION

The IGZ introduced the outcome indicator 'Percentage of patients with an unexpectedly long length of stay' for its supervision since 2010. It is based on the assumption that complications often prolong the patient's hospital stay. A higher percentage of patients with a UL-LOS compared to the national average, after a correction is made for case mix variations, may indicate shortcomings in the quality or safety of care delivered by the hospital. The indicator currently 'counts' all patients from whom the actual length of stay exceeds the expected duration by 50% or more. It needs to be studied whether this cut-off point can be set separately for each patient group. The indicator varies systematically between hospitals and is rather stable over time. It correlates with other outcome indicators, which could indicate the capacity to identify opportunities for improvement of patient safety.

ACKNOWLEDGMENTS

COMPETING INTERESTS

The authors declare that they have no competing interests.

FUNDING

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Annex 1. All Dutch hospital patients divided by length of stay (LOS) groups.

		Percentage of patients 2009	Percentage of patients 2010	Percentage of patients 2011
UL-LOS	observed LOS >100% longer than expected LOS	8,1%	7,9%	7,1%
	observed LOS 90-100% longer than expected LOS	0,9%	0,9%	0,8%
	observed LOS 80-90% longer than expected LOS	1,2%	1,0%	0,9%
	observed LOS 70-80% longer than expected LOS	1,1%	1,1%	1,0%
	observed LOS 60-70% longer than expected LOS	1,2%	1,3%	1,2%
	observed LOS 50-60% longer than expected LOS	1,6%	1,7%	1,6%
NON UL-LOS	observed LOS 40-50% longer than expected LOS	2,2%	2,0%	1,9%
	observed LOS 30-40% longer than expected LOS	2,1%	2,2%	2,0%
	observed LOS 20-30% longer than expected LOS	3,1%	2,9%	2,7%
	observed LOS 10-20% longer than expected LOS	3,2%	3,5%	3,4%
	observed LOS <10% longer than expected LOS	4,6%	5,0%	4,8%
	observed LOS ≤ expected LOS	68,6%	68,5%	70,8%
	deceased patients	2,1%	2,0%	1,8%
TOTAL		100,0%	100,0%	100,0%

FIGURE LEGENDS

- Figure 1: LOS = length of stay
 UL-LOS = unexpectedly long length of stay
- Figure 2: LMR = National Medical Registration
 HSMR = Hospital Standardised Mortality Rate
- Figure 3: I = 95% CI
 • = Admissions with UL-LOS 2011 (%)
- Figure 4: • = hospital
 UL-LOS = unexpectedly long length of stay

REFERENCES

1 Inspectie voor de Gezondheidszorg [Dutch Health Care Inspectorate]. *Basisset kwaliteitsindicatoren ziekenhuizen 2013*. Utrecht: IGZ; 2012.

2 Inspectie voor de Gezondheidszorg [Dutch Health Care Inspectorate]. *Basisset veiligheidsindicatoren ziekenhuizen 2010-2012*. Utrecht: IGZ; 2010.

3 Kwaliteitsindicatoren Zichtbare Zorg Ziekenhuizen. Available at: <http://www.zichtbarezorg.nl/page/Ziekenhuizen-en-ZBC-s/Kwaliteitsindicatoren#ZI>. Accessed 12/01, 2011.

4 Nolte E, Roland M, Guthrie S, et al. Preventing emergency readmissions to hospital, A scoping review. RAND Europe; 2012.

5 Lagoe RJ, Nanno DS, Luziani ME. Quantitative tools for addressing hospital readmissions. *BMC Res Notes* 2012;**5**:620,0500-5-620.

6 Goldfield NI, McCullough EC, Hughes JS, et al. Identifying potentially preventable readmissions. *Health Care Financ Rev* 2008;**30**:75-91.

7 Joynt KE, Jha AK. Thirty-day readmissions--truth and consequences. *N Engl J Med* 2012;**366**:1366-9.

8 Jarman B, Pieter D, van der Veen A, et al. The hospital standardised mortality ratio: a powerful tool for Dutch hospitals to assess their quality of care?. *Qual Saf Health Care* 2010;**19**:9-13.

9 Hoonhout LH, de Bruijne MC, Wagner C, et al. Nature, occurrence and consequences of medication-related adverse events during hospitalization: a retrospective chart review in the Netherlands. *Drug Saf* 2010;**33**:853-64.

10 Hoonhout LH, de Bruijne MC, Wagner C, et al. Direct medical costs of adverse events in Dutch hospitals. *BMC Health Serv Res* 2009;**9**:27.

-
- 11 Sari AB, Sheldon TA, Cracknell A, et al. Extent, nature and consequences of adverse events: results of a retrospective casenote review in a large NHS hospital. *Qual Saf Health Care* 2007;**16**:434-9.
- 12 Ehsani JP, Jackson T, Duckett SJ. The incidence and cost of adverse events in Victorian hospitals 2003-04. *Med J Aust* 2006;**184**:551-5.
- 13 Cho SH, Ketefian S, Barkauskas VH, et al. The effects of nurse staffing on adverse events, morbidity, mortality, and medical costs. *Nurs Res* 2003;**52**:71-9.
- 14 Camp M, Chang DC, Zhang Y, et al. Risk factors and outcomes for foreign body left during a procedure: analysis of 413 incidents after 1 946 831 operations in children. *Arch Surg* 2010;**145**:1085-90.
- 15 Lotfipour S, Kaku SK, Vaca FE, et al. Factors associated with complications in older adults with isolated blunt chest trauma. *West J Emerg Med* 2009;**10**:79-84.
- 16 Williams DJ, Olsen S, Crichton W, et al. Detection of adverse events in a Scottish hospital using a consensus-based methodology. *Scott Med J* 2008;**53**:26-30.
- 17 Kaushal R, Bates DW, Franz C, et al. Costs of adverse events in intensive care units. *Crit Care Med* 2007;**35**:2479-83.
- 18 Rice-Townsend S, Hall M, Jenkins KJ, et al. Analysis of adverse events in pediatric surgery using criteria validated from the adult population: justifying the need for pediatric-focused outcome measures. *J Pediatr Surg* 2010;**45**:1126-36.
- 19 Schioler T, Lipczak H, Pedersen BL, et al. Incidence of adverse events in hospitals. A retrospective study of medical records. *Ugeskr Laeger* 2001;**163**:5370-8.
- 20 Baker GR, Norton PG, Flintoft V, et al. The Canadian Adverse Events Study: the incidence of adverse events among hospital patients in Canada. *CMAJ* 2004;**170**:1678-86.

21 Davis P, Lay-Yee R, Briant R, et al. Adverse events in New Zealand public hospitals I: occurrence and impact. *N Z Med J* 2002;**115**:U271.

22 Lagoe R, Pernisi L, Luziani M, et al. Addressing hospital length of stay outlier patients: A community wide approach. *Advances in Bioscience and Biotechnology* 2014;**6**:188-96.

23 de Bruijne MC, Zegers M, Hoonhout LHF, et al. Onbedoelde schade in Nederlandse ziekenhuizen: dossieronderzoek van ziekenhuisopnames in 2004. Amsterdam: EMGO Instituut/VUmc en Nivel, Nederlands Instituut voor onderzoek van de gezondheidszorg; 2007.

24 Langelaan M, Baines RJ, Broekens MA, et al. Monitor Zorggerelateerde Schade 2008, dossieronderzoek in Nederlandse ziekenhuizen. NIVEL; 2010.

25 Zegers M, de Bruijne MC, Wagner C, et al. Adverse events and potentially preventable deaths in Dutch hospitals: results of a retrospective patient record review study. *Qual Saf Health Care* 2009;**18**:297-302.

26 Silber JH, Rosenbaum PR, Koziol LF, et al. Conditional Length of Stay. *Health Serv Res* 1999;**34**:349-63.

27 Aylin P, Bottle A, Majeed A. Use of administrative data or clinical databases as predictors of risk of death in hospital: comparison of models. *BMJ* 2007;**334**:1044.

28 Landelijke Medische Registratie (LMR). Available at: <http://www.dutchhospitaldata.nl/registraties/lmrlazr/Paginas/default.aspx>. Accessed 06/10, 2013.

29 Hart 't P, de Boer K, Draaisma D. On the length of stay in hospitals. Radboud University Nijmegen; 2012.

30 Cihangir S, Borghans I, Hekkert K, et al. A pilot study on record reviewing with a priori patient selection. *BMJ Open* 2013;**3**:10.1136/bmjopen.2013-003034. Print 2013.

31 Bottle A, Aylin P. Intelligent information: a national system for monitoring clinical performance. *Health Serv Res* 2008;**43**:10-31.

32 Bottle A, Middleton S, Kalkman CJ, et al. Global comparators project: international comparison of hospital outcomes using administrative data. *Health Serv Res* 2013;**48**:2081-100.


33 Marang-van de Mheen PJ, Lingsma HF, Middleton S, et al. Evaluation of quality of care using registry data: the interrelationship between length-of-stay, readmission and mortality and impact on hospital outcomes. *BMJ Qual Saf* 2014;**23**:350-1.

34 Website of more than 40 small and medium-sized Dutch hospitals. They take more than a third of all Dutch hospital care on their behalf. Available at: <http://www.saz-ziekenhuizen.nl>. Accessed 09-30, 2013.


35 Website with facts and figures of all (8) Dutch university hospitals. Available at: <http://www.nfu.nl/umc/feiten/>. Accessed 09-30, 2013.

36 Website of 28 Dutch tertiary teaching hospitals. Available at: <http://www.stz.nl/pagina/9-over-stz.html>. Accessed 09-30, 2013.


37 Rivard PE, Luther SL, Christiansen CL, et al. Using patient safety indicators to estimate the impact of potential adverse events on outcomes. *Med Care Res Rev* 2008;**65**:67-87.




90x44mm (300 x 300 DPI)



90x127mm (300 x 300 DPI)



127x90mm (300 x 300 DPI)



90x63mm (300 x 300 DPI)

Annex 1. All Dutch hospital patients divided by length of stay (LOS) groups.

		Percentage of patients 2009	Percentage of patients 2010	Percentage of patients 2011
UL-LOS	observed LOS >100% longer than expected LOS	8,1%	7,9%	7,1%
	observed LOS 90-100% longer than expected LOS	0,9%	0,9%	0,8%
	observed LOS 80-90% longer than expected LOS	1,2%	1,0%	0,9%
	observed LOS 70-80% longer than expected LOS	1,1%	1,1%	1,0%
	observed LOS 60-70% longer than expected LOS	1,2%	1,3%	1,2%
	observed LOS 50-60% longer than expected LOS	1,6%	1,7%	1,6%
NON UL-LOS	observed LOS 40-50% longer than expected LOS	2,2%	2,0%	1,9%
	observed LOS 30-40% longer than expected LOS	2,1%	2,2%	2,0%
	observed LOS 20-30% longer than expected LOS	3,1%	2,9%	2,7%
	observed LOS 10-20% longer than expected LOS	3,2%	3,5%	3,4%
	observed LOS <10% longer than expected LOS	4,6%	5,0%	4,8%
	observed LOS ≤ expected LOS	68,6%	68,5%	70,8%
	deceased patients	2,1%	2,0%	1,8%
TOTAL		100,0%	100,0%	100,0%

For peer review only